

# Ravulizumab-cwvz (Ultomiris)

Provider Order Form rev. 01/02/2024

## PATIENT INFORMATION

Referral Status:  New Referral  Updated Order  Order Renewal

Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_ Patient Phone: \_\_\_\_\_  
Patient Address: \_\_\_\_\_ Patient Email: \_\_\_\_\_  
Allergies: \_\_\_\_\_  NKDA Weight (lbs/kg): \_\_\_\_\_ Height (in/cm): \_\_\_\_\_  
Sex:  M /  F Date of Last Infusion: \_\_\_\_\_ Next Due Date: \_\_\_\_\_ Preferred Location: \_\_\_\_\_

## DIAGNOSIS (Please provide ICD-10 code in space provided)

Myasthenia Gravis (anti-acetylcholine receptor antibody positive): \_\_\_\_\_  
Other: \_\_\_\_\_ Description: \_\_\_\_\_

## REQUIRED INFORMATION

Must have meningococcal vaccine at least 2 weeks prior to infusions. If no vaccine, referring provider must provide 2 weeks of antibacterial drug prophylaxis

## THERAPY ADMINISTRATION & DOSING

Administer Ultomiris IV over 1 hour (**Choose one**):

**Weight 40-60kg:**

- Loading: 2400mg (in 24ml NS) at week 0, followed by 3000mg (in 30ml NS) at week 2
- Maintenance: 3000mg (in 30ml NS) every 8 weeks

**Weight 60-100kg:**

- Loading: 2700mg (in 27ml NS) at week 0, followed by 3300mg (in 33ml NS) at week 2
- Maintenance: 3300mg (in 33ml NS) every 8 weeks

**Weight 100kg or more:**

- Loading: 3000mg (in 30ml NS) at week 0, followed by 3600mg (in 36ml NS) at week 2
- Maintenance: 3600mg (in 36ml NS) every 8 weeks

**Switching from Eculizumab:** Administer loading dose 2 weeks after last dose of eculizumab followed by maintenance dose every 8 weeks

Monitor Patient for 60mins after every infusion

**DO NOT mix 300mg/30ml vials with other concentrations. OK to mix 300mg/3ml and 1100mg/11ml together if needed.**

## PRE-MEDICATION ORDERS

Tylenol  500mg /  650mg PO  
 Loratadine 10mg PO  
 Pepcid 20mg  PO /  IVP  
 Benadryl  25mg /  50mg  PO /  IVP  
 Solumedrol  40mg /  125mg IVP  
 Other: \_\_\_\_\_

## NURSING

Hold infusion and notify provider for:

- abnormal vital signs or signs/symptoms of infection or Meningitis
- New or worsening headache or altered mental status

Record vitals before infusion then every 30mins until patient discharges. If reactions occur, slow or stop infusion

Provide nursing care per Nursing Procedure, including Hypersensitivity Reaction Management Protocol and post-procedure observation.

## ADDITIONAL ORDERS

## PROVIDER INFORMATION

Preferred Contact Name: \_\_\_\_\_ Preferred Contact Email: \_\_\_\_\_  
Ordering Provider: \_\_\_\_\_ Provider NPI: \_\_\_\_\_  
Referring Practice Name: \_\_\_\_\_ Phone: \_\_\_\_\_ Fax: \_\_\_\_\_  
Practice Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip Code: \_\_\_\_\_

## REQUIRED DOCUMENTATION CHECKLIST (Additional documentation required for processing and insurance approval)

**Required Documentation:** Patient demos, copy of front and back of primary and secondary insurance, 2 most recent OVN including treatment failures or contraindications, EMG results, MRI results

**Required Labs:** AChR antibody, MuSK antibodies, CRP, ESR

Provider Name (*print*)

Provider Signature

Date

Order valid for one year unless otherwise indicated. IV solutions/diluents may be substituted as allowed per manufacturer's instructions as necessitated by product availability.