# Risankizumab-rzaa (Skyrizi IV)

Provider Order Form rev. 08/28/2024

# OI Infusion Services

PATIENT INFO	RMATION	<b>Referral Status:</b>	🗆 New R	eferral	Updated Ord	er 🛛 Order Renewal
Patient Name:			DOB:	DOB: Patient Pho		ne:
Patient Address:			Patient Email:			
Allergies:			□ NKDA	Weigh	t (lbs/kg):	Height (in/cm):
Sex: 🗆 M / 🗆 F	Date of Last Infusion:	Next Due Date	:	Pref	erred Location:	

## DIAGNOSIS (Please provide ICD-10 code in space provided)

Crohn	's Di	sease	(IV	dosing	only	):

Ulcerative Colitis (IV dosing only):

## **THERAPY ADMINISTRATION & DOSING**

☑ Only IV induction dosing will be provided. Subcutaneous dosing **WILL NOT** be provided

#### For Crohn's Disease:

□ Administer Risankizumab-rzaa (Skyrizi) 600mg IV over 1hour. Dilute in 100ml 0.9 NS or D5W

#### For Ulcerative Colitis:

□ Administer Risankizumab-rzaa (Skyrizi) 1200mg IV over 2 hours. Dilute in 250ml 0.9 NS or D5W

#### FREQUENCY (Choose one)

☑ Induction: week 0, week 4, and week 8

## **ADDITIONAL ORDERS**

## LABORATORY ORDERS

□ Bilirubin, LFTs at week 8 □ Other:

## **PRE-MEDICATION ORDERS**

□ Tylenol □ 500mg / □ 650mg PO

- □ Loratadine 10mg PO
- □ Pepcid 20mg □ PO / □ IVP

□ Benadryl □ 25mg / □ 50mg □ PO / □ IVP

□ Solumedrol □ 40mg / □ 125mg IVP

□ Other: \_

# NURSING

☑ Hold infusion and notify provider for:

- Signs or symptoms of illness/active infection or recent
- live vaccinationsElevated LFTs or bilirubin
- ☑ Provide nursing care per Nursing Procedure, including Hypersensitivity Reaction Management Protocol and post-

procedure observation

#### **PROVIDER INFORMATION**

Preferred Contact Name:	Pref	Preferred Contact Email:			
Ordering Provider:	Provider NPI:				
Referring Practice Name:	Phone:	Fax:			
Practice Address:	City:	State:	Zip Code:		

#### **REQUIRED DOCUMENTATION CHECKLIST (Additional documentation required for processing and insurance approval)**

**Required Documentation:** Patient demos, copy of front and back of primary and secondary insurance, 2 most recent OVN including treatment failures or contraindications with immunosuppressants, biologic agent and steroids, Colonoscopy **Required Labs:** TB, Hep B, CRP, ESR, LFTs and Bilirubin,

Provider Name (print)

**Provider Signature** 

Date

Order valid for one year unless otherwise indicated. IV solutions/diluents may be substituted as allowed per manufacturer's instructions as necessitated by product availability.