

Iron Infusion (Feraheme, Venofer, Monoferric, Injectafer)

Provider Order Form rev. 12/12/2024

PATIENT INFORMATION

Referral Status: New Referral Updated Order Order Renewal

Patient Name: _____ DOB: _____ Patient Phone: _____
Patient Address: _____ Patient Email: _____
Allergies: _____ NKDA Weight (lbs/kg): _____ Height (in/cm): _____
Sex: M / F Date of Last Infusion: _____ Next Due Date: _____ Preferred Location: _____

DIAGNOSIS (Select from list or provide ICD-10 code in space provided)

D50.0: iron deficiency secondary to blood loss D50.8: Other iron deficiency Anemia
 D63.0: Anemia in neoplastic disease D63.1: Anemia in CKD E83.10: Disorder of iron metabolism, unspecified
Other: _____ Description: _____

THERAPY ADMINISTRATION (Choose one)

Infuse iron product as required by patient's insurance.
List in order of preference: _____, _____, _____
 Infuse this product only (subject to prior authorization)

DOSING & FREQUENCY

Venofer Dose: 100mg, 200mg, 300mg IV. Mix 100mg and 200mg in 100ml NS and infuse over 15min. Mix 300mg in 250mg NS and infuse over 90min.

Venofer Frequency: (Choose one)

- every ___ days for ___ doses
 every ___ weeks for ___ doses

Feraheme Dose & Frequency:

administer 510mg IV x2 doses (Separated by 3-8 days). Mix in 100ml NS and infuse over 15-30 minutes.

Injectafer Dose & Frequency:

Pts over 50kg, administer 750mg IV on day 0 and day 7
 Pts under 50kg, administer 15mg/kg IV = _____ mg on day 0 and day 7
Mix in 250ml NS and infuse over 30 minutes.

Monoferric Dose & Frequency

Pts over 50kg, administer 1000mg IV over at least 20min as single dose. Dilute in 100ml NS
 Pts under 50kg, administer 20mg/kg IV = _____ mg over at least 20mins as single dose. Dilute to final concentration of 1mg/ml

PRE-MEDICATION ORDERS

Tylenol 500mg / 650mg PO
 Loratadine 10mg PO
 Pepcid 20mg PO / IVP
 Benadryl 25mg / 50mg PO / IVP
 Solumedrol 40mg / 125mg IVP
 Other: _____

NURSING

Hold infusion and notify provider for history of allergy to IV iron
 Monitor patient for hypersensitivity reaction for 30 minutes post infusion.
 Place patient in reclined or semi-reclined position.
 Use with caution in patients with hypotension (*feraheme/venofer*)
 Provide nursing care per Nursing Procedure, including Hypersensitivity Reaction Management Protocol and post-procedure observation

ADDITIONAL ORDERS

PROVIDER INFORMATION

Preferred Contact Name: _____ Preferred Contact Email: _____
Ordering Provider: _____ Provider NPI: _____
Referring Practice Name: _____ Phone: _____ Fax: _____
Practice Address: _____ City: _____ State: _____ Zip Code: _____

REQUIRED DOCUMENTATION CHECKLIST (Additional documentation required for processing and insurance approval)

Required Documentation: Patient demos, copy of front and back of primary and secondary insurance, 2 most recent OVN including treatment failures or contraindications with oral iron, Reason for anemia (by indication)

Required Labs: Kidney function, CBC, Ferritin, Iron, TIBC, Iron saturation, Iron within the last 4 weeks.

Provider Name (*print*) _____

Provider Signature _____

Date _____

Order valid for one year unless otherwise indicated. IV solutions/diluents may be substituted as allowed per manufacturer's instructions as necessitated by product availability.